Original Research Article

Prevalence and risk factors of allergic conjunctivitis in a semi urban set up

Manikala Sailaja¹, Muduthanpally Charani^{1,*}¹Dept. of Ophthalmology, Mallareddy Institute of Medical Sciences and Hospital, Hyderabad, Telangana, India

ARTICLE INFO

Article history:

Received 01-07-2020

Accepted 11-07-2020

Available online 30-03-2021

Keywords:

Allergic conjunctivitis

Allergic rhinitis

Prevalence

ABSTRACT

Background: Inflammation of the conjunctiva, which is caused due to exposure to allergens is called allergic conjunctivitis. Simple allergic conjunctivitis is fairly common and the symptoms are usually underrated and, most of the patients do not go to the hospital for treatment. There is a wide variation in the occurrence of allergic conjunctivitis in different geographical parts of the world.

Materials and Methods: 158 of the patients with allergic conjunctivitis were included in the study were found to have allergic conjunctivitis. Complete history including symptoms such as redness, itching, teary eyes etc. were taken in detail.

Results: 53.2% were in the primary school, while 37.3% were in the middle school, or between 6-10th class and 9.5% were in the 11th or 12th std. Exposure to dust was the main cause for allergic conjunctivitis as was seen in 32.3% of the cases. Smoke was the second most common cause with 22.8% affected, followed by Dye eyes in 13.9% of the cases. All the patients experienced itching in the eye, and were rubbing their eyes often. (98.7%). 86.1% had watery eyes with over Lacrimal secretion. Majority of the patients (81.6%) with allergic conjunctivitis had allergic rhinitis. 32.3% had asthma, and 13.3% had atopic dermatitis.

Conclusion: Since Allergic conjunctivitis is a mild disease with little morbidity, the patients fail to understand the importance of a regular eye checkup. This can be corrected with the help of health education to the students and the patients in the community.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Inflammation of the conjunctiva, which is caused due to exposure to allergens is called allergic conjunctivitis.¹⁻³ Some allergic conjunctivitis are more common and does not endanger the vision like seasonal allergic conjunctivitis, giant papillary conjunctivitis and perennial allergic conjunctivitis, while some threaten the vision and are less prevalent like atopic keratoconjunctivitis, and vernal keratoconjunctivitis.⁴ This type of conjunctivitis is usually Type I atopic hypersensitivity, which is IgE mediated, many times, with the involvement of cell mediated Th2.⁵

Simple allergic conjunctivitis is fairly common and the symptoms are usually underrated and, most of the patients do not go to the hospital for treatment. As a result, it is very

difficult to estimate the prevalence of this condition. It is however estimated that about 10-30% of the population are affected by this disorder. In United States around 40% of the people are affected by this disorder.⁶⁻⁸

Allergic conjunctivitis most frequently associated with allergic rhinitis. Other atopic diseases such as dust allergy, food allergy, eczema is also associated with allergic conjunctivitis.^{9,10}

The simple allergic conjunctivitis are acute, seasonal or perinneal. These are IgE mediated and results in the activation of the mast cells by the allergens and subsequent release of active substances like histamine and causing the dilatation of the blood vessels and irritation of nerve endings, resulting in to symptoms.¹¹ Mechanism of vernal kerato conjunctivitis is not very well understood, but is also said to be because of IgE and also T cell involvement. Atopic kerato conjunctivitis is expected to be due to IgE as

* Corresponding author.

E-mail address: sailajamanikala@gmail.com (M. Charani).

well as a delayed hypersensitivity reaction.^{12,13}

The mode of treatment is generally antihistamines, which provide fast relief. There is a wide variation in the occurrence of allergic conjunctivitis in different geographical parts of the world. Therefore, this study was conducted to see the prevalence of allergic conjunctivitis in this part of country.

2. Materials and Methods

This observational study was conducted in the department of ophthalmology at Mallareddy institute of medical sciences and hospital, Telangana for a period of three years during December 2016 to August 2019. Over 3000 children between the ages 5 to 16 years were surveyed at 3 different schools near our hospital. Out of them 1398 children had various vision problems and were asked to come to the hospital for further treatment. Out of these children, 158 of them were found to have allergic conjunctivitis. The nature of the study was explained to the patients and informed consent was taken from all of them. After clearance from the institutional ethical committee, demographic details were collected from all the patients. The nature and the duration of the conjunctivitis was noted. The socioeconomic status of the child, the nature of the family and its size was also noted.

All children whose parents refused to give consent or whose red eye was diagnosed not to be allergic conjunctivitis were excluded from the study.

Visual acuity was checked for all the patients using Snellen lettered chart as well as the tumbling letter E chart. A slit lamp biomicroscope was used to assess the anterior segment and the posterior segment was assessed using the indirect and the direct ophthalmoscopes. History of redness, itching, photophobia, tears secretion, mucinous secretions were taken down in detail.

3. Results

Out of the 158 children included in the study, 67 (42.4%) of them were females and 91 (57.4%) were males (Figure 1).

The most common age group which was affected with allergic conjunctivitis was between 5-7 years with 64 children affected (40.5%). 49 children between the ages 8-10 (31%), 26 (16.6%) between 11-13 and, 19 (12%) between 14-16 years were seen to be affected (Figure 2).

Most of the children were in their primary school, many of them in the 1st or 2nd standard. 84 (53.2%) were in the primary school, while 59 (37.3%) were in the middle school, or between 6-10th class and 15 (9.5%) were in the 11th or 12th std. Most of them were Hindus, which accounted for 78 (49.4%), 44 (29.8%) were muslims and 30 (19%) were Christians. 4 (2.5%) were Sikhs and 1 was a parsi and 1 was unable to say the religion. 86 (54.4%) were in a nuclear family while 72 (45.6%) were living in a joint family. 76

Fig. 1: Gender wise distribution

Fig. 2: Age wise distribution

(48%) had a pet like a dog, a cat or a rabbit at home or in the compound (Table 1).

Exposure to dust was the main cause for allergic conjunctivitis as was seen in 51 (32.3%) of the cases. Smoke was the second most common cause with 36 (22.8%) affected, followed by Dry eyes in 22 (13.9%) of the cases. 19 (12.1%) of the children had AC due to excessive exposure to sunlight and pollen affected 14 (8.9%) of the cases (Table 2).

All the patients experienced itching in the eye, and were rubbing their eyes often. (98.7%). 136 (86.1%) had watery eyes with over lacrimal secretion. Ropy discharge was seen in 13 (8.2%) of the cases and photophobia was experienced by 18 students (11.4%). 28 (17%) of the patients experienced gritty eyes, as if “sand was inside the eye” (Table 3).

Majority of the patients with allergic conjunctivitis had allergic rhinitis i.e 129 patients, amounting to 81.6%. 51 (32.3%) had asthma, and 21 (13.3%) had atopic dermatitis (Table 4).

Table 1: Demographic details

Details	Number	Percentage
Education:		
Primary	84	53.2%
Middle school	59	37.3%
High school/Secondary school	15	9.5%
Religion:		
Hindu	78	49.4%
Muslim	44	27.8%
Christian	30	19%
Sikh	4	2.5%
Others	2	1.3%
Family lifestyle		
Joint Family	72	45.6%
Nuclear family	86	54.4%
Size of family		
< 4 members	54	34.2%
4-7 members	59	37.3%
8-11 members	24	15.2%
12-15 members	19	12%
>15 members	2	1.3%
Pets	76	48%

Table 2: Causes of allergic conjunctivitis

Causes	Number	Percentage
Sunlight	19	12.1%
Smoke	36	22.8%
Dust	51	32.3%
Dry eye	22	13.9%
Pollen	14	8.9%
Dandruff	11	7%
Bacterial conjunctivitis	3	1.9%
Glaucoma	1	0.6%
Vit A deficiency	1	0.6%

Table 3: Symptoms

Itching	158	100%
Rubbing of eye	157	98.7%
Watery eyes	136	86.1%
Redness of eye	61	38.6%
Ropy discharge	13	8.2%
Photophobia	18	11.4%
Grittiness	28	17.7%
Clear mucinous discharge	11	7%

Table 4: Other allergic conditions

Allergic comorbidities	Number	Percentage
Allergic rhinitis	129	81.6%
Asthma	51	32.3%
Atopic dermatitis	21	13.3%

4. Discussion

Allergies are known to affect around 20% of the world's population.^{14,15} They are often familial. Eyes and nose are the most common sites for the allergens to attack and thus, one of the more common allergic reactions is allergic conjunctivitis. Some of the most common allergens are dust, pollen, mold spores, animal dander etc. this results in redness of the eyes and itchy. Most of the times the AC is mild, so the patients do not seek medical help and it goes unnoticed. Repeated reactions of the allergens cause it to become severe.^{15–18}

The prevalence of AC in a study by Wade et al. was 7.4%, while in the present study it was 5.2%.¹⁹ Another study from Tanzania by Hall et al. reported 35% of the children presenting with AC.²⁰ In a study in Ghana, a very high prevalence of 40% was observed,²¹ and in yet another study, by Abah et al. from Zaire, reported only 7.3% prevalence, similar to our study.²² In the present study, the predominant gender affected was male, but this was not found to be significant. In the study by Wade et al., females were more in number than males with AC, but like our study, this difference too was not significant.¹⁹ Uchio et al. also reported higher number of females with AC, while another study by Marback et al. reported more males than females.^{23,24}

40.5% of the children in our study belonged to the age group between 5-7 years and 31% were between 8-10 years. 12% of children between 16-14 years were affected. AC is said to occur in children above the age of 3 years and continues till puberty. It then resolves itself in most of the cases.²⁵

100% of the patients experienced itching in the eye, and were rubbing their eyes often, while 86% had excessive lacrimation. Ropy discharge was seen in 8.2%, while photophobia was seen in 11.4%. In a survey by Singh and Sinha, itching was the most common symptom corroborating our study. The other symptoms associated with this condition was redness, irritation, water from the eyes.¹⁹

The most common age group which was affected with allergic conjunctivitis was between 5-7 years with 64 children affected (40.5%). 49 children between the ages 8-10 (31%), 26 (16.6%) between 11-13 and 19 (12%) between 14-16 years were seen to be affected..

Exposure to dust was the main cause for allergic conjunctivitis as was seen in 51 (32.3%) of the cases. Smoke was the second most common cause with 36 (22.8%) affected, followed by dry eyes in 22 (13.9%) of the cases 19 (12.1%) of the children had AC due to excessive exposure to sunlight and pollen affected 14 (8.9%) of the cases. A study by Wade et al. reported 60.7% of allergic conjunctivitis in the dry season, the cause being chiefly pollen allergy.¹⁹

Allergic rhinitis was the most common comorbidity in the present study in more than 80% of the patients, followed

by Asthma in 50% of the patients. The other comorbidity that was associated with allergic conjunctivitis was Allergic dermatitis. Rasario et al. reported that most of the children with allergic conjunctivitis had allergic rhinitis, resulting in a lower quality of life.⁶ In a study by Singh and Sinha, the prevalence of AC was 30% of the total OPD in Ophthalmology clinics in North India. They found a variation in the season in their study. The said statistics was in the summer months, while in the others, it was far less, of only 10%.²⁶ A study by Gradman et al. in Denmark reported 42% of Ac to be associated with rhinitis, while 30% had eczema and 24% had asthma.⁹ In a study by Malu et al., the most common comorbid condition to be observed was refractive error in 15.4% of the cases.²⁵

5. Conclusion

In many of the cases ocular allergic reaction is overlooked as it is minor discomfort and does not lead to severe infections. As a result, patients do not come to the hospital and seek medical attention. Only when the symptoms become more severe, do they approach a doctor.

The best way to communicate to the people the importance of a regular eye checkup, is health education by the professionals working in the community. This would create an awareness among the people, so that they would take a timely action in treatment.

6. Source of Funding

None.

7. Conflict of Interest

None.

References

- Bielory L, Friedlaender MH. Allergic Conjunctivitis. *Immunol Allergy Clin North Am*. 2008;28(1):43–58. doi:10.1016/j.iac.2007.12.005.
- Ono SJ, Abelson MB. Allergic conjunctivitis: Update on pathophysiology and prospects for future treatment. *J Allergy Clin Immunol*. 2005;115(1):118–22. doi:10.1016/j.jaci.2004.10.042.
- Ortega BZ, Vargas MAR, Tsuchiya FMI, Navarro BER, Monge J. Allergic conjunctivitis in children. *Rev Alerg Mex*. 2007;54(2):41–53.
- Rosa ML, Lionetti E, Reibaldi M, Russo A, Longo A, Leonardi S, et al. Allergic conjunctivitis: a comprehensive review of the literature. *Ital J Pediatr*. 2013;39(1):18. doi:10.1186/1824-7288-39-18.
- Maurya RP, Singh VP, Singh MK, Bhushan P. Ocular Allergy: An Overview Written in Book: “Update on Respiratory Disease & Allergy. Mishra JK, editor. Varanasi: Tara Book;.
- Rasario N, Bielory L. Epidemiology of allergic conjunctivitis. *Curr Opin Allergy Clin Immunol*. 2011;11(5):471–6. doi:10.1097/aci.0b013e32834a9676.
- Singh K, Axelrod S, Bielory L. The epidemiology of ocular and nasal allergy in the United States, 1988-1994. *J Allergy Clin Immunol*. 2010;126(4):778–83. doi:10.1016/j.jaci.2010.06.050.
- Brozek G, Lawson J, Szumilas D, Zejda J. Increasing prevalence of asthma, respiratory symptoms, and allergic diseases: Four repeated surveys from 1993-2014. *Respir Med*. 2015;109(8):982–90. doi:10.1016/j.rmed.2015.05.010.
- Gradman J, Wolthers OD. Allergic conjunctivitis in children with asthma, rhinitis and eczema in a secondary outpatient clinic. *Pediatr Allergy Immunol*. 2006;17(7):524–6. doi:10.1111/j.1399-3038.2006.00429.x.
- Whitehorn TFB, Spengel JM. The link between allergies and eosinophilic esophagitis: implications for management strategies. *Expert Rev Clin Immunol*. 2010;6(1):101–9. doi:10.1586/eci.09.74.
- Bielory L, Friedlaender MH. Allergic Conjunctivitis. *Immunol Allergy Clin North Am*. 2008;28(1):43–58. doi:10.1016/j.iac.2007.12.005.
- Jordakieva G, Jarolim EJ. The impact of allergen exposure and specific immunotherapy on circulating blood cells in allergic rhinitis. *World Allergy Organ J*. 2018;11(1):19. doi:10.1186/s40413-018-0197-0.
- Li A, Li S, Ruan F, Jie Y. Atopic keratoconjunctivitis. *Medicine*. 2018;97(16):e0372. doi:10.1097/md.00000000000010372.
- Hesselmar B, Aberg B, Eriksson B, Aberg N. Allergic rhinoconjunctivitis, eczema, and sensitization in two areas with differing climates. *Pediatr Allergy Immunol*. 2001;12(4):208–15.
- Nathan R, Meltze E, Selner J, Storms W. Prevalence of allergic rhinitis in the United States. *Curr Opin Allergy Clin Immunol*. 1997;99(6):S808–14. doi:10.1016/s0091-6749(97)80040-1.
- Baig R, Ali AW, Ali T, Ali A, Shah MN, Sarfaraz A. Prevalence of allergic conjunctivitis in school children of Karachi. *J Pak Med Assoc*. 2010;60(5):371–3.
- Biswas J, Saha I, Das D, Bandyopadhyay S, Ray B, Biswas G. Ocular morbidity among children at a tertiary eye care hospital in Kolkata, West Bengal. *Indian J Public Health*. 2012;56(4):293–6. doi:10.4103/0019-557x.106418.
- Singla JP, Parashar Y. Prevalence of allergic rhinitis, allergic conjunctivitis and atopic dermatitis in children with recurrent wheeze. *Indian Pediatr*. 2009;46:262–4.
- Wade PD, Iwuora AN, Lopez L, Muhammad MA. Allergic conjunctivitis at sheikh zayed regional eye care center, gambia. *J Ophthalmic Vis Res*. 2012;7(1):24–8.
- Hall A, Shilio B. Vernal keratoconjunctivitis. *Community Eye Health J*. 2005;18:76–8.
- Baba I. The red eye-first aid at the primary level. *Community Eye Health J*. 2005;18:70–2.
- Abah ER, Oladigbolu KK, Samaila E, Gani-Ikilama A. Ocular disorders in children in Zaria children’s school. *Niger J Clin Pract*. 2011;14(4):473–6. doi:10.4103/1119-3077.91759.
- Uchio E, Kimura R, Migita H, Kozawa M, Kadonosono K. Demographic aspects of allergic ocular diseases and evaluation of new criteria for clinical assessment of ocular allergy. *Graefes Arch Clin Exp Ophthalmol*. 2008;246(2):291–6. doi:10.1007/s00417-007-0697-z.
- Marback PMF, de Freitas D, Junior AP, Junior AP, Junior RB. Epidemiology and clinical features of allergic conjunctivitis in a reference center. *Arq Bras Oftalmol*. 2007;70(2):312–6.
- Malu KN. Allergic conjunctivitis in Jos-Nigeria. *Niger Med Jo*. 2014;55(2):166–70. doi:10.4103/0300-1652.129664.
- Singh M, Sinha BP. Allergic Conjunctivitis: a practice pattern survey from North India. *Indian J Appl Res*. 2019;9(11):10–4.

Author biography

Manikala Sailaja, Assistant Professor

Muduthanpally Charani, Assistant Professor

Cite this article: Sailaja M, Charani M. Prevalence and risk factors of allergic conjunctivitis in a semi urban set up. *Indian J Clin Exp Ophthalmol* 2021;7(1):83-86.